Gulls

North Sea

The defintion of gulls that this study will use is: Any of numerous long-winged web-footed aquatic birds within the subfamily Larinae of the family Laridae (Gull, 2024). Gulls are in general highly opportunistic omnivores, but primarily feed on fish, marine invertebrates and small mammals (Calvino-Cancela, 2011;



Annett & Pierotti, 1999; Ewins et al., 1994). However, various species gulls also eat fruit and 'garbage' if that is a more abundant food source (Calvino-Cancela, 2011; Annett & Pierotti, 1999; Ewins et al., 1994). This study identified 12 species of gulls that can be found in the North Sea (Hamilton, 1981; Courtens & Stienen, 2006; Jongbloed et al., 2023). The most common species of gulls include: Herring gull (*Larus argentatus*), Lesser black-backed gull (*Larus fuscus*), and Greater black-backed gull (*Larus marinus*) (Hamilton, 1981; Courtens & Stienen, 2006; Jongbloed et al., 2023).

History/ Population trends

Not much is known of the origin of Gulls. However, in the last couple of decades an increase can be found on reduction of gull population of almost all species of gull, identified by this study. The most prominent cause of population decline, as suggested by Duijns et al., is mortality because of wind turbines at sea (2020). According to Duijns et al., especially large gulls are at risk of colliding with wind turbines and the numbers of collision victims from wind farms can be predicted using a collision-rate model (Duijns et al., 2020). These predicted collision numbers are modelled using turbine-related parameters (turbine size, rotor height, speed) and species-related parameters (bird density, flight height distribution, flight speed, percentage of time in flight, night activity, wingspan, size, avoidance rate) (Duijns et al., 2020).

Miscellaneous

- The largest species of gull in the world is the Greater black-backed gull (Olsen, 2018). The Greater black-backed gull reaches an average wingspan of 150cm to 170cm and an average weight of 1.5kg to 1.8kg (Olsen, 2018).
- Gulls are most commonly found in marine environments where freshwater is rare. In order to get enough to drink, gulls have evolved a specialized gland that can separate salt from water (Fänge et al., 1958). This allows gulls to drink seawater, instead of freshwater (Fänge et al., 1958).

Sources

- Annett, A., & Pierotti, R. (1999). LONG-TERM REPRODUCTIVE OUTPUT IN

 WESTERN GULLS:CONSEQUENCES OF ALTERNATE TACTICS IN DIET

 CHOICE. *Ecological Society of America*, 80(1), 288–297.

 https://esajournals.onlinelibrary.wiley.com/doi/epdf/10.1890/0012-9658%281999%29080%5B0288%3ALTROIW%5D2.0.CO%3B2
- Calvino-Cancela, M. (2011). Gulls (Laridae) as frugivores and seed dispersers. *Plant Ecology*, 212(7), 1149–1157. https://doi.org/10.1007/s11258-011-9894-2
- Courtens, W., & Stienen, W. M. (2006). Marine biological valuation of seabirds in the Belgian Part of the North Sea. *Instituut Voor Natuur- En Bosonderzoek*. https://www.vliz.be/imisdocs/publications/125701.pdf
- Duijns, S., Helberg, M., Verstraete, H., Stienen, E. W. M., & Fijn, R. C. (2020). *Origin of large gulls in the North Sea* (By Rijkswaterstaat Zee en Delta; No. 19–257).

 Rijkswaterstaat Zee en Delta. Retrieved November 22, 2024, from https://dlwqtxts1xzle7.cloudfront.net/88555314/19_0290_Origin_of_large_gulls_in_t he_North_Sea-libre.pdf?1657741461=&response-content-disposition=inline%3B+filename%3DOrigin_of_large_gulls_in_the_North_Sea_A.pd f&Expires=1732269233&Signature=Milak1qrDTYQmbYQjNmkZze7c6eNw7cWeu GxDkwpb04oA3Q-dXzD6qbkv4Au0IbtikCH9l7Hs2RuDLtaKX32lV3ug4MrGDttRdU7cIqFCNBpfNxar

dXzD6qbkv4Au01btikCH917Hs2RuDLtaKX32IV3ug4MrGDttRdU7cIqFCNBpfNxar X8rEsCEjdv0fx90vgZGFF3F71jEmygLxZ4VrdIZ0ncKGxdbMMeVVHOSgDDm9F y1ueTxopfNJxRELr2W4KzK-

bNNz4SdFA3D2xgcvWtZIOQjhsyjnqxptkajNkbCBjPUSIhc~RA30muYnlnaXZ5IJgz P4Ucs7zRRZhQpQEjAijCX~LckF774iTmcSTKBKPLqNTSXYmspOLNHBlpYd~V -CdIZjZzDwBzXdz8PYg__&Key-Pair-Id=APKAJLOHF5GGSLRBV4ZA

- Ewins, P. J., Weseloh, D. V., Groom, J. H., Dobos, R. Z., & Mineau, P. (1994). The diet of Herring Gulls (Larus argentatus) during winter and early spring on the lower Great Lakes. *Hydrobiologia*, 279–280(1), 39–55. https://doi.org/10.1007/bf00027839
- Fänge, R., Schmidt-Nielsen, K., & Osaki, H. (1958). THE SALT GLAND OF THE HERRING GULL. *Biological Bulletin*, 115(2), 162–171. https://doi.org/10.2307/1539022
- Gull. (2024). In *Merriam-Webster Dictionary*. Retrieved November 21, 2024, from https://www.merriam-webster.com/dictionary/gull
- Hamilton, N. (1981). A BIBLIOGRAPHICALLY-ANNOTATED CHECKLIST OF THE BIRDS OF SHETLAND. *MeRLEWOOD RESEARCH AND DEVELOPMENT*PAPER, 0308–3675. https://nora.nerc.ac.uk/id/eprint/6814/1/R%26D084.pdf
- Jongbloed, R., Tamis, J., Van Der Wal, J., De Vries, P., Grundlehner, A., & Piet, G. (2023).

 Quick scan of cumulative impacts on the North Sea biodiversity: With a focus on selected species in relation to future developments in offshore wind energy.

 https://library.wur.nl/WebQuery/wurpubs/623068
- Olsen, M. (2018). Gulls of the world. In *Google Books*. Bloomsbury Publishing Plc.

 Retrieved November 22, 2024, from

 https://books.google.nl/books?hl=nl&lr=&id=LZhPDwAAQBAJ&oi=fnd&pg=PP1&dq=gulls+AND+%22largest+species%22+OR+%22Biggest+wingspan%22&ots=aRAX14C8Je&sig=2yJoKqKmuXBXT1W9sjvsLsvEmnw&redir_esc=y#v=onepage&q=gulls%20AND%20%22largest%20species%22%20OR%20%22Biggest%20wingspan%22&f=false